- [0320] Clayton A et al., Analysis of antigen presenting cell derived exosomes, based on immuno-magnetic isolation and flow cytometry. J Immunol Methods. 2001; 247(1-2):163-74.
- [0321] Conde-Vancells J., Rodriguez-Suarez E., Gonzalez E., Berisa A., Gil D., Embade N., Valle M., Luka Z., Elortza F., Wagner C., Lu S. C., Mato J. M., Falcon-Perez M. (2010) Candidate biomarkers in exosome-like vesicles purified from rat and mouse urine samples. Proteomics Clin. Appl. 4, 416-425.
- [0322] Crescitelli R et al., Distinct RNA profiles in subpopulations of extracellular vesicles: apoptotic bodies, microvesicles and exosomes. J Extracell Vesicles. 2013 Sep. 12; 2.
- [0323] George C et al., Increased risk of respiratory diseases in adults with Type 1 and Type 2 diabetes. Diabetes Res Clin Pract. 2018 August; 142:46-55. doi: 10.1016/j. diabres.2018.05.029. Epub 2018 May 24.
- [0324] Gupta S1 et al., Biomarkers of Systemic Inflammation and Risk of Incident Hearing Loss. Ear Hear. 2019 July/August; 40(4):981-989. doi: 10.1097/AUD. 00000000000000678.
- [0325] Khaw K T et al., Association of hemoglobin Alc with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. Ann Intern Med. 2004 Sep. 21; 141(6):413-20. PMID: 15381514.
- [0326] Koga K., Matsumoto K., Akiyoshi T., Kubo M., Yamanaka N., Tasaki A., Nakashima H., Nakamura M., Kuroki S., Tanaka M., Katano M. (2005) Purification, characterization and biological significance of tumorderived exosomes. Anticancer Res. 25, 3703-3707.
- [0327] Kannagi R et al., Stage-specific embryonic antigens (SSEA-3 and -4) are epitopes of a unique globoseries ganglioside isolated from human teratocarcinoma cells. EMBO J. 1983; 2(12):2355-61.
- [0328] Kinzebach S and Bieback K. Expansion of Mesenchymal Stem/Stromal cells under xenogenic-free culture conditions. Adv Biochem Eng Biotechnol. 2013; 129:33-57.
- [0329] Mathias R A et al., Isolation of extracellular membranous vesicles for proteomic analysis. Methods Mol Biol. 2009; 528:227-42.
- [0330] Nathan D M et al (Diabetes Control and Complications Trial Research Group), The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. N Engl J Med. 1993 Sep. 30; 329(14):977-86.
- [0331] Nilsson J et al., (2009) Prostate cancer-derived urine exosomes: A novel approach to biomarkers for prostate cancer. Br. J. Cancer 100, 1603-1607
- [0332] Palmisano G et al., Characterization of membraneshed microvesicles from cytokine-stimulated-cells using proteomics strategies. Mol Cell Proteomics. 2012 August; 11(8):230-43. doi: 10.1074/mcp.M111.012732.
- [0333] Robinson J, Halliwell J A, Hayhurst J H, Flicek P, Parham P, Marsh S G E: The IPD and IMGT/HLA database: allele variant databases. Nucleic Acids Research. 2015 43:D423-431.
- [0334] Selvin E et al., Meta-analysis: glycosylated hemoglobin and cardiovascular disease in diabetes mellitus. Ann Intern Med. 2004 Sep. 21; 141(6):421-31.

- [0335] Sidney J, Peters B, Frahm N, Brander C, Sette A. HLA class I supertypes: a revised and updated classification. BMC Immunol. 2008 Jan. 22; 9:1. doi: 10.1186/1471-2172-9-1.
- [0336] Silva et al., "Evaluation of HLA matchmaker compatibility as predictor of graft survival and presence of anti-HLA antibodies," Transplantation Proceedings, vol. 42, no. 1, pp. 266-269, 2010.
- [0337] Umanath K et al., Update on Diabetic Nephropathy: Core Curriculum 2018. Am J Kidney Dis. 2018 June; 71(6):884-895. doi: 10.1053/j.ajkd.2017.10.026.
- [0338] Valenzuela and Reed, "Antibodies in transplantation: the effects of HLA and non-HLA antibody binding and mechanisms of injury," in Transplantation Immunology. Methods in Molecular Biology (Methods and Protocols), vol. 1034, pp. 41-70, Humana Press, Totowa, N.J., USA, 2013.

What is claimed is:

- 1. A method of ameliorating impaired glucose tolerance in a subject in need thereof, comprising: administering to the subject a pharmaceutical composition, comprising placental adherent stromal cells (ASC), thereby ameliorating impaired glucose tolerance.
- 2. The method of claim 1, wherein said subject is at least 60 years of age at the onset of treatment.
- 3. The method of claim 1, wherein said subject has a HbA1c value of at least 43.5 mmol/mol at the onset of treatment.
- **4**. The method of claim **1**, wherein said subject has a body mass index of at least 27.5 kg/m² at the onset of treatment.
- 5. A method of reducing systemic inflammation in a subject with impaired glucose tolerance in a subject in need thereof, comprising: administering to the subject a pharmaceutical composition, comprising placental adherent stromal cells (ASC), thereby reducing systemic inflammation in a subject with impaired glucose tolerance.
 - 6-8. (canceled)
- **9**. The method of claim **1**, wherein said ASC have been incubated on a 3D culture substrate in a bioreactor.
- 10. The method of claim 9, further comprising the subsequent step of harvesting said ASC by removing said ASC from said 3D culture apparatus.
- 11. The method of claim 9, wherein said ASC have been incubated on a 2D adherent-cell culture substrate, prior to incubation in said 3D culture apparatus.
 - 12. (canceled)
- 13. The method of claim 9, wherein said 3D culture substrate comprises a synthetic adherent material formed as a fibrous matrix, wherein said synthetic adherent material is selected from the group consisting of a polyester, a polypropylene, a polyalkylene, a polyfluorochloroethylene, a polyvinyl chloride, a polystyrene, a polysulfone, a cellulose acetate, a glass fiber, a ceramic particle, a poly-L-lactic acid, and an inert metal fiber.
 - 14-16. (canceled)
- 17. The method of claim 1, wherein said administering comprises:
 - a. administering to the subject a first pharmaceutical composition, comprising allogeneic placental ASC from a first donor; and